

University of Groningen

## (S)-3,3-Dimethyl-1,2,4-butanetriol as Ligand for Titanium Catalysed Asymmetric Silylcyanation

Callant, Dominique; Stanssens, Dirk; Vries, Johannes G. de

*Published in:*  
Tetrahedron%3A Asymmetry

*DOI:*  
[10.1016/S0957-4166\(00\)82333-2](https://doi.org/10.1016/S0957-4166(00)82333-2)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1993

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Callant, D., Stanssens, D., & Vries, J. G. D. (1993). (S)-3,3-Dimethyl-1,2,4-butanetriol as Ligand for Titanium Catalysed Asymmetric Silylcyanation. *Tetrahedron%3A Asymmetry*, 4(2).  
[https://doi.org/10.1016/S0957-4166\(00\)82333-2](https://doi.org/10.1016/S0957-4166(00)82333-2)

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



$\text{LiAlH}_4$  reduction of D-pantolactone (1) followed by hydrolysis with  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$  gave the triol 2<sup>13</sup>, which after distillation and careful drying over molecular sieves was reacted with titanium tetra-isopropoxide in  $\text{CH}_2\text{Cl}_2$  during 4 hours to give the catalyst 3 as a solid.  $^1\text{H-NMR}$  of 3 confirmed the 1:1 ratio of triol-isopropoxide but showed mainly broad absorptions, characteristic of oligomeric titanium complexes. Curiously, the remaining isopropoxide peak appeared as a sharp singlet.

Unfortunately, no reaction occurred when 3 was added in stoichiometrical amount to a mixture of HCN and benzaldehyde in toluene. On the other hand, 3 proved to be an excellent catalyst in the asymmetric trimethylsilylation of aromatic aldehydes (Scheme 2.).



Scheme 2. *i*, TMSCN, 3, solvent; *ii*, 1N  $\text{H}_2\text{SO}_4$


Reaction of trimethylsilyl cyanide (TMSCN) with benzaldehyde at  $0^\circ\text{C}$  catalysed by an equivalent amount of 3 proceeded smoothly in a number of solvents (Table, entries 1-7) giving S-benzaldehyde cyanohydrin (4) after acidic hydrolysis in excellent yields with e.e.'s<sup>14</sup> up to 66%. Addition of activated 4Å molecular sieves gave a slight improvement in e.e. which could be augmented by lowering the temperature to  $-20^\circ\text{C}$ , leading to a maximum e.e. of 76%. No reaction occurred when the temperature was lowered further to  $-65^\circ\text{C}$ . Using 3 in catalytical amounts (10 mol%) led to a much slower reaction and had a detrimental effect upon the e.e. which decreased with conversion (entry 12). Curiously, addition of two equivalents of catalyst also gave a much lower e.e. The effect of the TMSCN concentration on the e.e. was probed by adding the reagent during one hour using the conditions of entry 9. This resulted in a lowered yield of 85% and an e.e. of 70%.

By adding hexane or pentane to the reaction mixture after the reaction is complete the catalyst could be precipitated and recovered by filtration. The recovered catalyst could be reused and under the conditions of entry 9 gave product with 71% e.e.. Since no apparent decomposition of the catalyst occurs under the reaction conditions it is not clear why the catalytical reaction gives a much lower e.e.. It is possible that the catalyst is converted into another oligomeric form after one turnover, which is reversed after precipitation with hexane. We are currently investigating this and other questions relating to the mechanism.

The scope of the use of this catalyst in asymmetric silylcyanation of other aldehydes remains to be determined. However, when 3-hydroxybenzaldehyde was subjected to the conditions of entry 9 no reaction took place. On the other hand, silylcyanation of the trimethylsilyl protected derivative 5 took place in good yield (Scheme 3.). The cyanohydrin was obtained

as a solid material of unknown configuration and had an e.e of 30%.

There is no mention in the literature of asymmetric trimethyl silylation of ketones. Attempted silylcyanation of p-isobutyl-acetophenone under a number of different conditions using 3 as catalyst were unsuccessful.



Scheme 3. *i*, 3, TMSCN, CH<sub>2</sub>Cl<sub>2</sub>; *ii*, 1N H<sub>2</sub>SO<sub>4</sub>

Table. Trimethylsilylation of benzaldehyde

Entry	Conditions <sup>a</sup>	Reaction time (h)	Yield <sup>b</sup> (%)	e.e (%)
1.	Toluene	1	92	60
2.	CH <sub>2</sub> Cl <sub>2</sub>	1	95	65
3.	Diethyl ether	1.25	93	54
4.	THF	1	92	66
5.	Hexane <sup>c</sup>	1.5	90	52
6.	Diphenyl ether	1	95	56
7.	Diisopropyl ether	1.5	90	40
8.	CH <sub>2</sub> Cl <sub>2</sub> , 4Å MS <sup>d</sup>	1	98	72
9.	CH <sub>2</sub> Cl <sub>2</sub> , 4Å MS, -20°C	2	92	76
10.	THF, -65°C		0	
11.	THF, 4Å MS, 0.1 eq. 3	12	70	10
12.	CH <sub>2</sub> Cl <sub>2</sub> , 4Å MS, 0.1 eq. 3	1		48
		3		44
		5	60	38
13.	CH <sub>2</sub> Cl <sub>2</sub> , 4Å MS, 2 eq. 3	1	90	40
14.	CH <sub>2</sub> Cl <sub>2</sub> , 4Å MS, 2 eq. TMSCN	1	95	66

a. Reaction performed at 0°C with equimolar amounts of benzaldehyde, TMSCN and 3, unless otherwise indicated

b. Determined by <sup>1</sup>H NMR

c. Catalyst remains largely undissolved

d. 100mg of 4Å MS/mmol of 3

## References

1. W.R. Jackson, H.A. Jacobs, G. Jayatilake, B.R. Matthews and K.G. Watson, *Aust. J. Chem.*, 1990, **43**, 2045.
2. C. Kruse, J. Brussee, *Spec. Chem.*, 1992, **12**, 184.
3. W. Becker, H. Freund and E. Pfeil, *Angew. Chem.*, 1965, **77**, 1139.
4. U. Niedermeyer and M.-R. Kula, *Angew. Chem.*, 1990, **102**, 423.
5. K. Tanaka, A. Mori and S. Inoue, *J. Org. Chem.*, 1990, **55**, 181.
6. K. Inoue, M. Matsumoto, S. Takahashi, T. Ohashi, and K. Watanabe, European Patent Application 271 868, 1988.
7. H. Minamikawa, S. Hayakawa, T. Yamada, N. Iwasawa and K. Narasaka, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 4379.
8. M. Hayashi, T. Matsuda and N. Oguni, *J. Chem. Soc. Chem. Commun.*, 1990, 1364.
9. a. A. Mori, H. Nitta, M. Kudo and S. Inoue, *Tetrahedron Lett.*, 1991, **32**, 4433. b. H. Nitta, D. Yu, M. Kudo, A. Mori and S. Inoue, *J. Am. Chem. Soc.*, 1992, **114**, 7969.
10. A. Mori, H. Ohno, H. Nitta, K. Tanaka and S. Inoue, *Syn. Lett.*, 1991, 563.
11. S. Kobayashi, Y. Tsuchiya and T. Mukaiyama, *Chem Lett.* 1991, 541.
12. For a zirconium catalyst based on a chiral amino-triol see: W.A. Nugent, *J. Am. Chem. Soc.*, 1992, **114**, 2768.
13. P. Lavallée, R. Ruel, L. Grenier and M. Bissonnette, *Tetrahedron Lett.*, 1986, **27**, 679.
14. Enantiomeric excess was determined using HPLC on a Daicel chiral OD column (n-Hexane/2-propanol 9:1 as mobile phase)